

Unusual patterns of I-131 contamination

Mehmet OZGUVEN, Seyfettin ILGAN, Nuri ARSLAN, A. Ozgur KARACALIOGLU,
Dogangun YUKSEL and Sabri DUNDAR

Department of Nuclear Medicine, Gulhane Military Medical Academy and Medical Faculty, Ankara, Turkey

Whole body imaging with radioiodine can detect functioning metastases, which can often be effectively treated with appropriate amounts of radioiodine. Non-physiologic I-131 uptake detected on images is usually interpreted as suggesting functioning thyroid metastases. However, extra-thyroidal I-131 accumulation does not always imply thyroid cancer metastases and has been reported on many occasions, including various non-thyroidal neoplasms, and contamination by body secretions. In order to avoid unnecessary therapeutic interventions it is extremely important to properly distinguish false-positive sites of I-131 localization. Three patients with unusual radioiodine contamination patterns, either presented for the first time or rarely presented in the existing literature, were reported. Reported cases consist of contamination in hair (due to styling hair with sputum), contamination in neck (due to drooling during sleep) and, contaminated chewing gum. False positive contamination sources were clarified by careful examination of patients and further images when necessary.

Key words: I-131 scanning, thyroid cancer, contamination

INTRODUCTION

A LARGE BODY of evidence indicates that radioiodine (I-131) is of value in the management of well-differentiated thyroid cancer. Whole body imaging (WBI) with radioiodine can detect functioning metastases, which can often be effectively treated with proper amounts of radioiodine.¹

Following thyroid ablation, physiologic activity is expected in the genitourinary tract, gastrointestinal tract, salivary glands, oropharynx, nasopharynx and esophagus.^{2,3} Non-physiologic I-131 uptake detected on images usually is interpreted as suggesting functioning thyroid metastases. However, extra-thyroidal I-131 accumulation does not always imply thyroid cancer metastases and has been reported in many circumstances,^{3–21} including various non-thyroidal neoplasms,^{22–26} and contamination by body secretions.^{2,27–31} There is little doubt that patients

with distant metastases that concentrate I-131 should receive I-131 therapy. In order to avoid unnecessary therapeutic interventions it is extremely important to properly distinguish false-positive sites of I-131 localization. We report here several examples of unusual contamination patterns on I-131 WBI in patients with well-differentiated thyroid carcinoma.

Standard approach to I-131 WBI for thyroid carcinoma in our department is to withdraw thyroid hormone replacement at least 4–6 weeks prior to the study, and to establish a serum thyroid stimulating hormone (TSH) level of $>30 \mu\text{IU/ml}$. I-131 WBI is performed using a large-field-of-view gamma camera (GE 400 ACT/STARCAM, Milwaukee, WI, USA) equipped with a high-energy parallel hole collimator. I-131 whole-body and appropriate static images were obtained at 24 and 48 hours after p.o. administration of 185 MBq diagnostic dose of I-131 or seven days after the administration of therapeutic doses of I-131.

CASE REPORT

Case I: Saliva Contamination of Hair

A 23-year-old male was referred to our center for I-131 therapy six weeks after total thyroidectomy and regional

Received October 29, 2003, revision accepted January 9 2004.

For reprint contact: Seyfettin Ilgan, M.D., GATA Medical Faculty, Department of Nuclear Medicine, 06018 Etlik-Ankara, TURKEY.

E-mail: silgan@gata.edu.tr

lymph node resection for papillary carcinoma. I-131 whole-body image after 5,550 MBq of I-131 therapy demonstrated multiple focal areas of uptake in the neck and thoracic regions suggesting nodal and lung metastases. Additionally diffuse accumulation of tracer surrounding the scalp was also noted (Fig. 1). TSH and thyroglobulin (Tg) levels at the time of imaging were $>75 \mu\text{IU/ml}$ and 16.8 ng/ml , respectively. Further inquiry revealed that he was used to style his hair using his sputum. The contamination disappeared after washing the hair.

Case II: Saliva Contamination in Neck

A 27-year-old male with follicular carcinoma undergone

routine follow-up scan five years after surgery and I-131 treatment. Diagnostic radioiodine WBI showed I-131 accumulation at the right lower neck with increasing intensity on two consecutive days (Fig. 2). TSH and Tg levels at the time of imaging were $>75 \mu\text{IU/ml}$ and 5.6 ng/ml , respectively. Activity persisted despite repeated washing of the skin. Finally it was recognized that contamination was due to sialorrhea (drooling) during sleep. Since the patient preferentially sleeps in the same position, contamination increasingly accumulated at the same location.

Case III: Radioactive Chewing Gum

A 57-year-old female who had undergone near-total thyroidectomy for papillary thyroid carcinoma was treated with 5,550 MBq of I-131. Post-therapeutic I-131 WBI demonstrated thyroid bed activity and diffuse hepatic uptake. Additionally, prominent focal tracer accumulation was observed in the right mandibular region, which is atypical for thyroid cancer metastases (Fig. 3). TSH and Tg levels at the time of imaging were $>75 \mu\text{IU/ml}$ and 10.8 ng/ml , respectively. On further investigation, the patient revealed that she had a chewing gum in her right buccal region, and repeated scan after the removal of chewing gum showed no activity in that region.

DISCUSSION

I-131 WBI is a widely accepted method for detecting functioning metastases of well-differentiated thyroid carcinoma, and non-physiologic uptake is often an indication to administer large amounts of radioiodine.¹ However, extra-thyroidal I-131 accumulation does not always denote thyroid cancer metastases and has been reported in




Fig. 1 (A) Anterior and posterior whole body images show multiple focal areas of uptake in the neck and thoracic region suggesting nodal and lung metastases beside diffuse accumulation of tracer surrounding the head. Anterior (B) and left lateral (C) static images of the head clearly demonstrate contamination of the hair.




Fig. 2 Anterior (A) and posterior (B) whole-body images show intense accumulation of I-131 at the right lower neck (*arrow*) due to saliva contamination. Contamination was due to drooling and persisted despite repeated washing of the skin.




Fig. 3 (A) Anterior static image of neck after therapeutic dose of I-131 showed intense focal uptake in the right mandibular region (*arrow*), beside thyroid bed activity (markers show mentum and sternal notch). On inquiry patient revealed that a chewing gum had been present in her right buccal area during imaging. After removal of the gum, activity disappeared (marker shows sternal notch) (B).

many circumstances including various non-thyroidal neoplasms, inflammations and contamination by body secretions and exudates.³⁻³⁰

Physiologic secretions of radioiodine from oropharynx, salivary glands, sweat glands, stomach and genitourinary tract are potential causes of misleading images. Radioiodine accumulation in the scalp due to excessive perspiration in a patient with a wig has been reported before.¹⁸ Though our case was similar to that with the

radioactive wig, contamination was in the hair and the underlying mechanism was a quite bizarre habit i.e. styling hair with sputum. This case of artificial accumulation reinforces the need to take into account patients' odd habits and fully inform them about potential contamination sources.

Although one can assume that drooling during night could be a potential cause of I-131 contamination no such case has been described in literature to our best knowledge. Since drooling continues overnight, contamination might be extensive and difficult to clean probably due to partial dermal absorption. Necessary precautions should also be taken against possible contamination of linen and pillows with I-131. Beside other potential uncommon saliva contamination types,²⁷⁻³⁰ drooling should be questioned in cases of suspected contamination.

Using hard sour candy or chewing gum to promote salivary secretions during radioiodine therapy is a common measure to reduce the radiation dose to salivary glands.³¹ Contaminated gums might mimic metastasis or dental pathologies and be a potential error source. Patients should be asked to change gum frequently and not to use it during imaging.

Although radioiodine has a very specific uptake mechanism in thyroid there are many conditions causing extrathyroidal I-131 accumulation other than thyroid cancer metastases. Nuclear medicine physicians should be aware of possible sources of contamination and false positive uptake patterns to prevent unnecessary I-131 treatments.

REFERENCES

1. Hurley JR, Becker DV. Treatment of thyroid carcinoma with radioiodine. In: *Diagnostic Nuclear Medicine*, Gottschalk A (ed). Baltimore; Williams & Wilkins, 1988: 792-814.
2. Camponovo EJ, Goyer PF, Silverman ED, Kistler AM, Yudt WM. Axillary iodine-131 accumulation due to perspiration. *Clin Nucl Med* 1989; 14: 762-763.
3. Wu SY, Brown T, Milne N, Egbert R, Kabok A, Lyons KP, et al. Iodine 131 total body scan—extrathyroidal uptake of radioiodine. *Semin Nucl Med* 1986; 16: 82-84.
4. Sutter CW, Benedicto GM, Stadalnik RC. False-positive results of I-131 whole-body scans in patients with thyroid cancer. *Semin Nucl Med* 1995; 15: 279-282.
5. Greenler DP, Klein HA. The scope of false-positive iodine-131 images for thyroid carcinoma. *Clin Nucl Med* 1989; 14: 111-117.
6. Kolla IS, Alazraki NP, Watts NB. Sialadenitis mimicking metastatic thyroid carcinoma. *Clin Nucl Med* 1989; 14: 564-566.
7. Geatti O, Shapiro B, Orsolon PG, Mirolo R, Di Donna A. An unusual false-positive scan in a patient with pericardial effusion. *Clin Nucl Med* 1994; 19: 678-682.
8. Bakheet SM, Hammami MM. False-positive radioiodine whole-body scan in thyroid cancer patients due to unrelated pathology. *Clin Nucl Med* 1994; 19: 325-329.
9. Hoschl R, Choy DH, Gandevia B. Iodine-131 uptake in

- inflammatory lung disease: A potential pitfall in treatment of thyroid carcinoma. *J Nucl Med* 1988; 29: 701–706.
10. Schneider JA, Divgi CR, Scott AM, Macapinlac HA, Sonenberg M, Goldsmith SJ, et al. Hiatal hernia on whole-body radioiodine survey mimicking metastatic thyroid cancer. *Clin Nucl Med* 1993; 18: 751–753.
 11. Ceccarelli C, Pacini F, Lippi F, Pinchera A. An unusual case of a false-positive iodine-131 whole-body scan in a patient with papillary cancer. *Clin Nucl Med* 1988; 13: 192–193.
 12. Caplan RH, Gundersen GA, Abellera RM, Kiskin WA. Uptake of iodine-131 by a Meckel's diverticulum mimicking metastatic thyroid cancer. *Clin Nucl Med* 1987; 12: 760–762.
 13. Brachman MB, Rothman BJ, Ramanna L, Tanasescu DE, Adelberg H, Waxman AD. False-positive iodine-131 body scan caused by a large renal cyst. *Clin Nucl Med* 1988; 13: 416–418.
 14. Achong DM, Oates E, Lee SL, Doherty FJ. Gallbladder visualization during post-therapy iodine-131 imaging of thyroid carcinoma. *J Nucl Med* 1991; 32: 2275–2277.
 15. Ziessman HA, Bahar H, Fahey FH, Dubianski V. Hepatic visualization on iodine-131 whole-body thyroid cancer scans. *J Nucl Med* 1987; 28: 1408–1411.
 16. Otsuka N, Fukunaga M, Morita K, Ono S, Nagai K, Katagiri M, et al. Iodine-131 uptake in a patient with thyroid cancer and rheumatoid arthritis during acupuncture treatment. *Clin Nucl Med* 1990; 15: 29–31.
 17. Norby EH, Neutze J, Nostrand DV, Burman KD, Warren RW. Nasal radioiodine activity: A prospective study of frequency, intensity, and pattern. *J Nucl Med* 1990; 31: 52–54.
 18. Abdel-Dayem H, Halker K, Sayed ME. The radioactive wig in iodine-131 whole body imaging. *Clin Nucl Med* 1985; 9: 454–455.
 19. Bakheet SM, Hammami MM. False-positive thyroid cancer metastasis on whole-body radioiodine scanning due to retained radioactivity in the oesophagus. *Eur J Nucl Med* 1993; 20: 415–419.
 20. Dugue JJ, Miguel MB, Ruiz E, Castillo L, Claver M, Rubio MJ, et al. False-positive I-131 whole-body scan in follicular thyroid carcinoma caused by frontal sinus mucocele. *Clin Nucl Med* 2000; 25: 137–138.
 21. Morgan R, Cote M. Abnormal uptake of I-131 mimicking salivary gland uptake in a patient with diffuse dental disease. *Clin Nucl Med* 2000; 25: 314–315.
 22. Kim EE, Pjura G, Gobuty A, Verani R. I-131 uptake in a benign serous cystadenoma of the ovary. *Eur J Nucl Med* 1984; 9: 433–435.
 23. Haubold-Reuter BG, Landolt U, Schulthess GK. Bronchogenic carcinoma mimicking metastatic thyroid carcinoma. *J Nucl Med* 1993; 34: 809–811.
 24. Lakshmanan M, Reynolds JC, Vecchio SD, Merino MJ, Norton JA, Robbins J. Pelvic radioiodine uptake in a rectal wall teratoma after thyroidectomy for papillary carcinoma. *J Nucl Med* 1992; 33: 1848–1850.
 25. Wang PW, Chen HY, Li CH, Chen WJ. Uptake of I-131 by an abdominal neurilemmoma mimicking metastatic thyroid carcinoma. *Clin Nucl Med* 1993; 18: 964–966.
 26. Ilgan S, Narin Y, Arslan N, Aksu A, Bayhan H. Warthin's tumor and I-131 body scan. *Clin Nucl Med* 1999; 24: 721–722.
 27. Gritters LS, Wissing J, Gross MD, Shapiro B. Extensive salivary contamination due to concurrent use of chewing tobacco during I-131 radioablative therapy. *Clin Nucl Med* 1993; 18: 115–117.
 28. Ain KB, Shih WJ. False-positive I-131 uptake at a tracheostomy site discernment with Tl-201 imaging. *Clin Nucl Med* 1994; 19: 619–621.
 29. Park HM, Tarver RD, Schauwecker DS, Burt R. Spurious thyroid cancer metastasis: Saliva contamination artifact in high dose iodine-131 metastases survey. *J Nucl Med* 1986; 27: 634–636.
 30. Roman MR, Larcos G, Devadas M. Lick your lips: a case of salivary contamination after I-131 therapy for thyroid carcinoma. *Clin Nucl Med* 2002; 27: 462.
 31. Harbert JC. Radioiodine Therapy of Differentiated Thyroid Carcinoma. In: *Nuclear Medicine Therapy*. Harbert JC (ed), New York; Thieme Medical Publishers, 1987: 72.