

Lung Scintigraphy with ^{131}I -MAA and Computer Processing of RI Image

H. YOSHIOKA, M. HASEGAWA and I. IWASAKI

Second Department of Internal Medicine, Okayama University

Medical School, Okayama

Lung scintigraphy by ^{131}I -MAA has been studied routinely as one of the lung function test. But it is difficult to evaluate quantitatively local blood flow of lung and dynamic blood flow.

To study this we used PHO/GAMMA scintillation camera, Tosbac 40 minicomputer, teletypewriter, videotaperecorder and diversing collimeter.

The locations, size and degree of diminished distributions in the pulmonary circulation had been made clear by smoothing the lung scintigraphy.

By profile curves of any axis of X or Y distribution of blood flow had been made clear.

RI changes with time any given region of lung and heart could be obtained.

Pulmonary Scintigraphic Abnormalities in Aortitis Syndrome; their Incidence and their Relation to Pulmonary Tuberculosis

M. MORI, T. TAKEDA, I. ITO, T. SHIRAISHI, S. KOIKE, Y. SASAKI,

M. IIO, and S. MURAO

Second Department of Internal Medicine, University of Tokyo, Tokyo

It has been well established that pulmonary arteries may be also involved in aortitis syndrome (Takayasu's arteritis). It is also known that the incidence of tuberculosis is high in this disease and that pulmonary perfusion scans are often abnormal in patients with known tuberculosis. Therefore we have at least two different causes by which pulmonary perfusion scan may become abnormal in this disease. The purpose of this study is to know the incidence of abnormal pulmonary perfusion scans in aortitis syndrome and to decide whether these abnormalities are related to pulmonary tuberculosis or aortitis syndrome per se.

Pulmonary perfusion scans were obtained from 38 cases of aortitis syndrome. Clinical histories and chest films were also carefully reviewed in each case. The diagnosis being confirmed by histories, physical examinations and finally by angiographic studies. ^{131}I -MAA about 0.3 mCi

was used for the lung scan and the injections were made in supine position.

Scans were abnormal in 28 out of 38 cases (73%). History of tuberculosis was obtained in 12, out of which scan was abnormal in 10, on the other hand, out of 26 patients without history of tuberculosis, scan was abnormal in 18. X-ray findings indicative of old tuberculous scars were observed in 13, out of which scan was abnormal in 11, while out of 25 patients without tuberculous lesions, scan was abnormal in 17. Though the incidence of abnormal pulmonary perfusion scans is slightly higher in patients with history of tuberculosis or chest film findings of old tuberculous scars, the difference was statistically not significant. There are, however, at least five cases in whom scintigraphic abnormalities were definitely related to old tuberculosis.

It is concluded that the incidence of abnormal perfusion scan is about 70% in aortitis syndrome