

mentioned, 4 patients with congenital heart disease, 3 patients of postcardiac surgery, 2 normal volunteers). Both early and delayed LAO30° views were used to evaluate changes in focal ^{201}Tl accumulation according to time in rest.

Twelve out of 40 patients showed right ventricular wall delineation, and nine out of 40 patients showed change in focal activity (usually area of decreased activity) between early and delayed LAO30° views. All patients other than infarction who showed area of decreased activity included apex. This was thought to be due to the fact that apex was periphery in coronary circulation. In 12 patients of right ventricular wall delineation, 10

had congestive heart failure, and in 30 patients without CHF 28 did not show right ventricular wall. Overall acuity of right ventricular wall delineation was 95%. In 6 patients who showed increased focal activity in delayed scan, 5 showed no organic change, and in 3 patients who showed decreased focal activity, 2 showed presence of organic change. Overall acuity in terms of organic change was 78%. With 6 view images all parts of myocardium was able to delineate tangentially and localization of any abnormal myocardial areas were possible. No acute adverse reaction due to toxicity of ^{201}Tl was detected in any of 40 patients.

Tl-201 Scintigraphy for Myocardial Imaging

M. KONDO*, A. KUBO*, F. KINOSHITA*, I. KITAGAWA*, K. OIKAWA* and H. YAMAZAKI**

**Department of Radiology School of Medicine, Keio University*

***Department of Internal Medicine*

TI-201 scintigraphy was performed in several disorders. Starting 10 minutes after intravenous injection of 2.0 mci of TI-201, scintiphotos were obtained in five different views (anterior, left lateral, 30, 45 and 60 degree left anterior oblique). Images were obtained with a Toshiba gamma-camera GCA-202. The energy spectrum used was 75 keV \pm 30%.

Studies were performed in 14 patients with acute or myocardia infarction, 6 with angina pectoris, 5 with cardiomyopathy, 7 with congenital heart disease, 6 with mitral stenosis and 5 with other disease.

Scintiphotos showed a defect in 6 of 8 patients with acute myocardial infarction and in 5 of 6 patients with old myocardial infarction. There was

a good agreement between electrocardiographic and scintigraphic location of the infarction. Scintiphotos showed no defects in patients with angina pectoris at rest.

Scintigraphy was also performed in patients with pressure or volume overload of the right ventricle such as ASD, PS and Tetralogy of Fallot. The right ventricle was fairly well visualized in those cases. Since with TI-201 the normal right ventricle is barely or not visualized, good visualization of the right ventricle per se may indicate the presence of the pressure or volume overload of the right ventricle. Also, in patients with heart failure or cardiomyopathy, scintigraphy will provide objective criteria of hypertrophy and/or dilatation of the right ventricle.

Myocardial Imaging in Idiopathic Cardiomyopathy Using Thallium-201

M. HAMADA*, T. NAKANO*, H. TAKEZAWA*, Y. OHIZUMI**, H. MAEDA**,
T. NAKAGAWA** and N. YAMAGUCHI**

**The First Department of Internal Medicine, Mie University School of Medicine, Tsu, Japan*

***Department of Radiology, Mie University School of Medicine, Tsu, Japan*

The purpose of this presentation is to evaluate the clinical validity of myocardial imaging using i.v. injection of Thallium-201 (^{201}Tl) in patient

with idiopathic cardiomyopathy.

23 cases of idiopathic cardiomyopathy (ICM), including 8 cases of hypertrophic obstructive type