

interpretation of the values obtained by RIA.
Possible relation of anti-T₃ antibody in Case 5

to latent hypothyroid state of this patient was also discussed.

Comparison of TSH Receptor Assays

Nobuaki KUZUYA*, S. C. CHIU*, H. IKEDA*, H. UCHIMURA*, K. ITO** and S. NAGATAKI**

**The Third Department of Internal Medicine, Faculty of Medicine, University of Tokyo*

***Ito Hospital, Tokyo*

There have been reported two representative methods for TSH receptor assays by Amir et al. (J.B.C., 1973) and by Smith et al. (FEBS Letters, 1974) after Manley (J. Endocr., 1974). However, these two methods are quite different from each other in both assay conditions and results of assays on the sensitivity to cold TSH and on the effects of IgGs from patients with Graves' disease. In order to observe what factor was the most responsible for these discrepancies between the two methods, this study was performed using the preparations prepared as described in the original reports from the same materials (e.g. receptor, tracer, cold hormone, IgG). Under the original condition, the displacement of ¹²⁵I-TSH by the cold TSH was as sensitive as that in the original report in each assay. Many IgGs from the patients with Graves' disease showed stronger displacements than those from normals in Smith's

method. However, both IgGs displaced the ¹²⁵I-TSH to the same extents in Amir's method despite of the use of same IgG preparations. Any alterations of incubation temperature, incubation time and pH resulted in decrease in binding and sensitivity in each assay. The degradation of ¹²⁵I-TSH during incubation or the amount of solubilized receptors in the incubation medium was negligible in both methods. Substitution of one receptor for another did not influence the results in each assay. Similarly, substitution of Smith's tracer for Amir's did not change the sensitivity or IgG's effect in Amir's assay. These results suggest that either methods to prepare receptors or degradations of labelled TSH are not responsible for the differences between the two assays. The combination of purified tracer and incubation conditions seems to be important to increase the sensitivity of TSH receptor assays.

Studies on the Radioreceptor Assay of TSH

—TBII in Patients Treated with Radioiodide for Hyperthyroidism—

K. ENDO*, K. KASAGI*, J. KONISHI*, T. OKUNO*, T. MORI**, K. KUMA*** and K. TORIZUKA*

**Department of Radiology and Nuclear Medicine, Kyoto University School of Medicine*

***Kobe Central City Hospital,*

****Kuma Hospital*

By using the radioreceptor assay of TSH, some IgG from patients with Graves' disease have been shown to inhibit the binding of labelled TSH to its receptor sites. These IgG, called TSH-binding inhibitor immunoglobulins (TBII), were detected in 60% of untreated Graves' patients. In this study TBII were measured in 51 patients who had been treated with radioiodide (¹³¹I, 4–10 mCi) 4 to 17 years before.

The incidence of TBII was 20%. In still thyro-

toxic patients (10 cases) TBII were detected more frequently (80%) and LATS activity was positive in 20%. However, the incidence of TBII in hypothyroid (12 cases) or euthyroid (29 cases) patients were very low of 5%. Furthermore the activities of TBII in these patients were not so potent as in thyrotoxic patients. This result may indicate that the measurement of TBII in patients treated with radioiodide is useful for checking the results of treatment.