

RECENT ADVANCE IN MOTILIN AND hPP RESEARCH

Kozaburo Mori, Yutaka Seino, Shigeru Matsukura,
Takuo Fujita, Zen Itoh*

Third Division, School of Medicine, Kobe University,

*Department of Surgery, Gumma University.

We studied the relationship between plasma motilin levels and gastric motor activity in conscious dogs and healthy subjects. Plasma motilin levels were measured by sensitive radioimmunoassay while gastric motor activity was recorded by means of chronically implanted force transducer in dogs and intragastric balloon tubing in man.

When the gastric contractile activity was in the interdigestive state, plasma motilin levels were always elevated in all dogs, furthermore it fluctuated in complete association with hunger contractions during the interdigestive state. Ingestion of food lowered the plasma motilin concentration and this low levels of motilin lasted as long as the gastric motor activity was in the digestive pattern. In human study, the similar findings were obtained between fasting motilin levels and gastric motor activity, furthermore gastric borborygmus closely associated with high motilin concentration in the fasting state.

Human pancreatic polypeptide (hPP) is newly recognized pancreatic hormone described Chance et al. Using radioimmunoassay, we studied fasting plasma levels of hPP and responses to food stimuli in healthy subjects and patients of chronic pancreatitis. Basal level of hPP in young control subjects was 50 ± 5 pg/ml, although older subjects showed high level of hPP according to each decade of age. There was a significant correlation of basal hPP and age. After ingestion of food plasma hPP increased promptly and showed biphasic secretory pattern in healthy subjects, these postprandial high level was sustained for 5 to 6 hours. Impaired release of hPP to food stimuli in patients with chronic pancreatitis were observed. So to study the hPP secretion to food stimuli is very useful and noninvasive tool for pancreatic exocrine and endocrine function.

RADIOIMMUNOASSAYS OF NEUROPEPTIDES

Shigeru Matsukura and Hiroki Yoshimi

Third Division, Department of Medicine, Kobe
University School of Medicine, Kobe.

Recently, it was reported that many bioactive substances were present in the central nervous system of several species including man. These substances are considered to function to convey information between the adjacent neurons (e.g. nor-epinephrine, dopamine and acetylcholine) or to modify, stimulate or inhibit, chemical neurotransmission (e.g. neuropeptides such as enkephalins, β -endorphin, LH-RH and TRH), which are called neurotransmitters and neuromodulators respectively. For the measurements of these compounds, relatively non-specific and insensitive bioassays have been used so far. However, recently, very specific and sensitive radioimmunoassays of these compounds, especially neuropeptides, have been developed and applied for analyses of these substances in the brain, the visceral organs, the cerebrospinal fluid and the blood.

We have developed a radioimmunoassay for β -endorphin using an anti-serum obtained in a rabbit immunized with β -endorphin obtained in the crude porcine ACTH preparations (Organon). The minimal detectable amount was 5 pg per tube. The antiserum used reacted with β_3 -LPH as equally as β -endorphin on a molar base but showed negligible cross-reactivity with other peptides including α -endorphin β -MSH, α -MSH and enkephalins, indicating that this antiserum was directed to 17-31 amino acid sequence of β -endorphin. Using this assay we have confirmed the presence of β -endorphin in the pituitary extracts of human, rat and cow in addition to β -LPH after fractionations by Sephadex G-100 column. Furthermore, we have measured the content of β -endorphin in the various brain regions and several visceral organs of the Japanese monkeys. The interpeduncular nucleus had the highest content of β -endorphin activity except for the pituitary, among the brain regions analyzed, followed by the habenula and the hypothalamus. Considerably higher amounts were found in the preoptic areas, the substantia nigra, the pallidum, the superior and the inferior colliculi, the olfactory tract, the corpus callosum, the pyramis and the ventral as well as the dorsal roots of the spinal cord. In contrast, no significant amount of immunoreactive β -endorphin was detected in the visceral organs such as the