

98

CLINICAL EVALUATION OF CARCINO-
EMBRYONIC ANTIGEN AND TISSUE
POLYPEPTIDE ANTIGEN IN PLEURAL
EFFUSION, ASCITES, AND BILE
K. Nakada, T. Tsukamoto, K. Fujimori,
T. Shougase, K. Ito, M. Furudate and
H. Omote*. Hokkaido University
School of Medicine, Sapporo.
*Hokkaido University Hospital Section
of Central Radiology, Sapporo.

Carcinoembryonic Antigen (CEA)
and Tissue Polypeptide Antigen (TPA)
levels in pleural effusion, ascites,
and bile were measured by radio-
immunoassay technique to evaluate
clinical value of these two tumor
associated antigens in the body
fluids. The body fluids were obtained
from 64 patients with malignant or
benign disease. Reproducibilities, and
recovery tests of the body fluids
proved to be less reliable than those
of sera, especially in bile. There was
no significant correlation between CEA
levels and TPA levels in the body
fluids. TPA levels in the body fluids
were so high both in malignant and
benign diseases that it was difficult
to set cut-off value like in serum
levels, but it was suggested measure
of TPA levels in the body fluids may
be useful in monitoring therapeutic
effects or progress of the disease.

99

CORRELATION BETWEEN SERUM CA19-9 LEVELS
AND LEWIS PHENOTYPE IN NORMAL SUBJECTS.
H. Sakahara, K. Endo, K. Nakajima,
T. Nakashima, M. Koizumi, H. Ohta, and
K. Torizuka. Kyoto University, Kyoto.

CA19-9 is a very useful tumor marker
for pancreatic carcinoma and other
malignant tumors. Since CA19-9 is a
sialylated Lewis A (Le^a) antigen, serum
CA19-9 concentrations have been considered
to be related to Lewis blood group
antigens. We examined the relationship
between serum CA19-9 levels and Lewis
phenotype in normal healthy subjects.

Serum CA19-9 level was 40.8 ± 19.3 U/ml
in Le(a+b-) group (N=10) and 11.4 ± 3.0 U/ml
in Le(a-b+) group (N=20). On the other
hand, CA19-9 was almost undetectable in
Le(a-b-) group (N=10). Individuals with
Le(a-b-) phenotype lack an enzyme that
catalyzes the synthesis of the common
sugar sequence of Lewis antigens and
CA19-9. As a consequence, it is considered
that they can not synthesize CA19-9.
Furthermore, when sera or their IgG
fraction obtained from some Le(a-b-) subjects
were added to the assay system, the
recovery of CA19-9 was very poor. Polyclonal
antisera against Le^a or Le^b also caused
similar results in recovery studies. Some
Le(a-b-) subjects seemed to have anti-Lewis
antibody which would cross-react with
CA19-9.

These results indicated a close
correlation between CA19-9 and Lewis blood
group antigens.

100

Ganglioside pattern in liver tissues and
distribution of Ca 19-9 on thin-layer chro-
matogram
M. Tanno, H. Yamada, J. Nagashima, A. Yamagata,
H. Nishino and K. Chiba. Tokyo Metropolitan
Geriatric Hospital, Tokyo.

Gangliosides were isolated from human li-
ver tissues with various liver diseases
and specific gangliosides which are capa-
ble to bind Ca19-9 antibody were identi-
fied on one and two-dimensional thin-layer
chromatography using the method of auto-
radiography.

Ganglioside pattern of normal tissues was
the predominance of GM3 and other minor
components on thin-layer chromatogram.
While, ganglioside patterns of hepatocellu-
lar carcinoma and certain type of liver
cirrhosis were found to be complicated as
compared with those of normal liver. These
characteristic features were marked increa-
sed amount of GM2 and several unidentifi-
ed gangliosides. On the study of autoradio-
graphy, there were two strong bands which
had activity of 123-I Ca19-9 in normal ti-
ssue on chromatogram. These bands had only
negligible amounts of ganglioside compone-
nt in the normal tissue.

Thus, these findings suggest that normal
tissue as well as hepatocellular tissue
have a ganglioside component designated as
sialylated lacto-fucopentaose II, Ca19-9
and its metabolic pathway.

101

EVALUATION OF SERUM CA19-9 IN PATIENTS WITH
PANCREATIC AND OTHER GASTROINTESTINAL
MALIGNANCIES.

T. Sasazawa, M. Ogata, T. Tani, Y. Kato.
The Research Institute of Tuberculosis
Hospital, Kiyose.

Although many diagnostic methods have
been developed, Pancreatic disease is dif-
ficult to diagnose correctly.

Recently, CA19-9 was detected as new
gastrointestinal tumor-associated antigen,
especially as pancreatic cancer.

We have tested serum CA19-9 of 120
patients with gastrointestinal diseases in
our hospital during last one year, and
compared the positive rate of each tumor
markers.

The reference range of 20 normal adults
was less than 5.5 U/ml to 21.5 U/ml ($8.7 \pm$
 6.2 U/ml).

When serum CA19-9 level higher than 30
U/ml were regarded as positive, it was
found predominantly in sera of gastro-
intestinal malignancies, especially in
pancreatic cancer.

We thought that serum CA19-9 was the
most reliable tumor marker with high sensi-
tivity and specificity for diagnosis of
pancreatic and biliary tract cancer.